





Responsible RestartOhio

Consumer, Retail & Services







Employees

Mandatory

Ensure min 6 ft between people, if not possible, install barriers

- Employees must perform daily symptom assessment*
- · Require employees to stay home if symptomatic
- Face coverings must be worn at all times
- Require regular handwashing
- Place hand sanitizers in high-contact locations
- Clean high-touch items after each use (e.g. carts, baskets)

Recommended Best Practices

Group employees by shift to reduce exposure

Customers & Guests

Ensure minimum 6 ft between people

- Specify hours for at-risk populations (e.g. elderly)
- Place hand sanitizers in high-contact locations
- Ask customers and guests not to enter if symptomatic
- Face coverings to be worn while shopping or visiting
- Stagger entry of customers and guest

- Health questionnaire for symptoms at entry point
- Provide face coverings upon entry
- Where possible, accept customers by appointment only
- Increase availability for curb-side pickup
- Consider suspending return policies

Physical Spaces

Ensure minimum 6 ft between people, if not possible, install barriers

- Post social distancing signage & disinfect high-contact surfaces hourly
- Clean merchandise before stocking if possible
- Establish maximum capacity (e.g. 50% of fire code)
- Discontinue self-service food stations, product samples
- Food courts remain closed

- Close once a week for deep cleaning
- Maximize available checkout space to promote social distancing (e.g., space customer lines with floor markers, use alternate registers)
- Use contact-less payments where possible
- Increase capacity for delivery and curb-side pickup

Confirmed Cases

- Immediately isolate and seek medical care for any individual who develops symptoms while at work
- Contact the local health district about suspected cases or exposures
- · Shutdown shop/floor for deep sanitation if possible
- Work with local health department to identify potentially infected or exposed individuals to help facilitate effective contact tracing/ notifications
- Once testing is readily available, test all suspected infections or exposures
- Following testing, contact local health department to initiate appropriate care and tracing

^{*}Daily symptom assessments should include taking your temperature with a thermometer and monitoring for fever. Also watch for coughing or trouble breathing.